$\sim$	1	Ω	4	a
C	1	<b>2</b>	4	J

(Pages: 3)

Name.....

Reg. No.....

# SIXTH SEMESTER (CUCBCSS-UG) DEGREE EXAMINATION, MARCH 2021

## Mathematics

MAT 6B 12-NUMBER THEORY AND LINEAR ALGEBRA

lime: Three Hours

Maximum: 120 Marks

## Section A

Answer all questions.

Each question carries 1 mark.

- 1. State Division algorithm.
- 2. Find gcd(272,1479).
- 3. Show that the Diophantine equation 14x + 35y = 651 has an integer solution.
- 4. Define e-prime. Give an example of a prime number.
- 5. Find the last two digits of the number 99°
- 6. State Fermat's theorem.
- 7. Find  $\tau(180)$ .
- 8. Define subspace of a vector space.
- 9. Give a basis for  $Mat_{2\times 2}(\mathbb{R})$ .
- 10. What do you mean by a Linear Transformation?
- Give an injective linear map which is not surjective.
- 12. State the Dimension theorem.

 $(12 \times 1 = 12 \text{ marks})$ 

#### Section B

Answer at least eight questions. Each question carries 6 marks. All questions can be attended. Overall Ceiling 48.

- 13 Show that the expression  $\frac{a(a^2+2)}{3}$  is an integer for al.  $a \ge 1$ .
- 14. If a|bc, with gcd(a, b) = 1, prove that a|c.

Turn over

- . . .
- 15. Use the Euclidean Algorithm to obtain integers x and y satisfying gcd(119, 272) = 119x + 2
- 16. Prove that the number  $\sqrt{5}$  is irrational.
- 17. Find the canonical form of 2093.
- 18. If  $p_n$  is the  $n^{th}$  prime number, prove that  $p_n \le 2^{2n-1}$ .
- 19. Find the remainder obtained upon dividing the sum 1! + 2! + 3! + ... + 99! + 100! by 12.
- 20. Let  $P(x) = \sum_{k=0}^{m} c_k x^k$  be a polynomial function of x with integral coefficients  $c_k$ . If  $a \equiv b(x_0)$  prove that  $P(a) \equiv P(b) \pmod{n}$ .
- 21. If  $(n-1)! \equiv -1 \pmod{n}$ , then prove that n must be prime.
- 22. Prove that every plane through the origin is a subspace of  $\mathbb{R}^3$ .
- 23. Show that  $\{(1, 1, 0, 0), (-1, -1, 1, 2), (1, -1, 1, 3), (0, 1, -1, -3)\}$  is a basis of  $\mathbb{R}^4$ .
- 24. Let S be a subset of the vector space V. Prove that S is basis if and only if S is a minimal span set.
- 25. Let  $f: \mathbb{R}^4 \to \mathbb{R}^3$  be a linear map given by f(a, b, c, d) = (a + b, b c, a + d). Find Im f and a for Im f.
- 26. Prove that if the linear mapping  $f: V \longrightarrow W$  is injective and  $\{v_1, v_2, ..., v_n\}$  is a linearly indepensable of V then  $\{f(v_1), f(v_2), ..., f(v_n)\}$  is a linearly independent subset of W.

 $(8 \times 6 = 48 \, \text{m})$ 

## Section C

Answer at least **five** questions. Each question carries 9 marks. All questions can be attended. Overall Ceiling 45.

- 27. Let a and b be integers, not both zero. Prove that a and b are relatively prime if and only if exist integers x and y such that 1 = ax + by.
- 28. Prove that if k > 0, then gcd(ka, kb) = kgcd(a, b).
- 29. State and Prove Fundamental Theorem of Arithmetic.
- 30. Prove that the system of linear congruences  $ax + by \equiv r \pmod{n}$ ;  $cx + dy \equiv s \pmod{n}$  has a use solution modulo n whenever gcd(ad bc, n) = 1.
- 31. If n is an odd pseudoprime, prove that  $M_n = 2^n 1$  is a larger one.

- 32. Let V be a vector space that is spanned by the finite set  $G = \{v_1, v_2, v_3, ..., v_n\}$ . If  $I = \{w_1, w_2, ..., w_m\}$  is linearly independent subset of V then show that  $m \le n$ .
- 33. Let V be finite dimensional vector space. If G is a finite spanning set of V and if I is a linearly independent subset of V such that  $I \subseteq G$  then prove that there is a basis B of V such that  $I \subset B \subseteq G$ .
- S4. Prove that a linear mapping is completely and uniquely determined by its action on basis.
- 35. If  $f: \mathbb{R}^3 \to \mathbb{R}^2$  is a linear map such that f(1, 1, 0) = (1, 2), f(1, 0, 1) = (0, 0), f(0, 1, 1) = (2, 1), then find f(x, y, z) for all  $(x, y, z) \in \mathbb{R}^3$ .

 $(5 \times 9 = 45 \text{ marks})$ 

#### Section D

Answer any one question.

The question carries 15 marks.

- 36. a) Let a and b be integers, not both zero. For positive integer d, prove that d = gcd(a, b) if and only if
  - (i) d|a and d|b
  - (ii) Whenever c|a and c|b, then c|d.
  - b) A customer bought a dozen pieces of fruit, apples and oranges, for \$1.32. If an apple costs 3 cents more than an orange and more apples than oranges were purchased, how many pieces of each kind were bought.
- 37. State and prove Chinese Remainder Theorem.
- 38. Let V and W be vector spaces each of dimension n over a field F. If  $f: V \longrightarrow W$  is linear then prove that the following statements are equivalent:
  - (i) f is injective;
  - (ii) f is surjective;
  - (iii) f is bijective;
  - (iv) f carries bases to bases.

 $(1 \times 15 = 15 \text{ marks})$