D 100636

(Pages: 2)

Reg. No....

SIXTH SEMESTER U.G. (CBCSS-UG) DEGREE EXAMINATION, MARCH 2024

Physics/Applied Physics

PHY6B12/APH 6B 12—NUCLEAR PHYSICS AND PARTICLE PHYSICS

(2019 Admission onwards)

Time: Two Hours

Maximum: 60 Marks

The symbols used in this question paper have their usual meanings.

Section A - Short Answer type

Answer all questions in two or three sentences, each correct answer carries a maximum of 2 marks.

- 1. Explain the correction in the binding energy formula obtained using liquid drop model.
- 2. Why do heavy nuclei have more neutrons than protons?
- 3. Write a short note on radio isotope production in nuclear reaction.
- 4. What is the strange behavior of kaons and hyperons?
- 5. What is the working principle of an intersecting beam accelerator?
- 6. Give the list of leptons. Mention the charge of each particle.
- 7. What do you mean by quantum chromodynamics?
- 8. In general, would you expect fission fragment to decay by positive or negative beta decay? Why?
- 9. Comment on the property of nuclear force.
- 10. List some similarities and difference between the properties of photons and neutrinos.
- 11. Explain the working of semiconductor counters.
- 12. Explain why a fusion reactor requires a high particle density, a high temperature and a long confinement time?

(Ceiling 20 marks)

Section B - Paragraph / Problem type

Answer all questions in a paragraph of about half a page to one page, each correct answer carries a maximum of 5 marks.

- 13. Consider a single helium nucleus formed by the fusion of two deuterium nuclei. Mass of $_1\mathrm{H}^2 = 2.014102u$; mass of $_2\mathrm{He}^4 = 4.002604u$. Find out the energy released in fusion.
- 14. Distinguish between fission and fusion reactions. Explain the fusion process in stars.

Turn over

471925

 p^{I}

- 15. Determine whether the following reactions are allowed or forbidden.
 - $p+p \rightarrow n^+ + p + \pi^+$.
 - $p + \pi^- \to \pi^0 + n.$ (ii)
 - (iii) $e^+e^+ \rightarrow \mu^+ + \pi^-$
 - 16. Discuss briefly low energy reaction kinematics.
 - The disintegration constant λ of a radioactive element is 0.00231 per day. Calculate its λ and average life.
 - 18. Discuss the Quark model.
 - 19. A reactor is developing energy at the rate of 3000kW. How many atoms of U^{235} undergo figs (Ceiling 30 second?

Section C - Essay type

Essays - Answer in about two pages, any one question. Answer carries 10 marks.

- 20. Using a neat diagram explain the working principle of Van de Graaff electrostatic genera
- 21. Explain the different elementary particle quantum numbers and their conservation la examples. $(1 \times 10 = 10)$